



Precisely target pests with **RETALIATE-BN**

Our RetaliatE-BN series of products is a specific range of Beneficial Nematode species.

These species, known as Entomopathogenic Nematodes or EPNs, have a life cycle comprised of the egg stage, four juvenile stages, and the adult stage. These EPNs have a free-living third stage juvenile (J3), termed a dauer juvenile, that is the infective stage and therefore called an IJ. These IJs harbor a symbiotic bacterium that plays an essential role in subsequent stages of the life cycle.

EPNs mortality to insects is a 3-step process:

1

FIRSTLY,

the nematode must migrate, searching for the target host.

2

SECONDLY,

upon contact with the host, it must penetrate through natural body openings, such as the anus, mouth and/or cuticle.

3

FINALLY,

the nematode-bacteria complex should be able to overcome the insect's immune system and multiply, producing new generations of IJs.



The nematodes infect insects and inject the bacteria into their hemolymph (circulatory fluid). Therefore, the bacteria multiply and produce a range of low-molecular-weight compounds and insecticidal proteins. Insect usually dies within 48-72 hours and serves the nematodes as a nutrient source and habitat for reproduction. After acquisition of new endosymbionts, the nematodes leave their host and search for other prey.

RETALIATE-BN/Sf

TARGETS: THRIPS, FUNGUS GNATS



Nematode Species

Steinernema feltiae

Symbiotic Bacteria

Xenorhabdus bovienii /

nematophilus

Foraging Characteristics

Uses both ambusher and cruiser means of finding hosts

Temperature Range

Active at temperatures of 55 - 82F

Primary Pests Controlled

Thrips (*Franlinothrips* sp.)

Fungus Gnats (*Bradysia* sp.)

Fruit Flies (*Drasophylla*)

Tray Concentration

1 tray contains 1.5 Billion juvenile (IJ)

S. feltiae OR 1 tray contains 500

Million juvenile (IJ) *S. feltiae* (2 trays per sleeve)

Dosage Recommendation:

1.5 Billion Trays

Early season soil drench: 1 tray per 2-4 acres

Foliar applications: twice weekly to twice monthly depending on pest infestation:

Light Rate: 1 tray treats 6-10 acres.

Heavy Rate: 1 tray treats 3-6 acres.

Dosage Recommendation:

500 Million Trays

Early season soil drench: 1 tray per 0.5-1.5 acres

Foliar applications: twice weekly to twice monthly depending on pest infestation:

Light Rate: 1 tray treats 2-3.5 acres.

Heavy Rate: 1 tray treats 1-2 acres.

Shelf Life

4 weeks. Unopened container. Keep refrigerated.

Other pests controlled:

Cabbage Maggot (*Delia radicum*)

Onion Maggot (*Delia antiqua*)

Crane Flies (*Dipetera* sp.)

Shore Flies (*Scatella stagnalis*)

**Dosages vary depending on pest species, infestation levels, and environmental factors. Consult your Account representative for pest specific dosages.*

DIRECTIONS

- Follow directions for crop and nematode species dosages.
- Pre-irrigate soil if necessary to achieve uniform soil moisture and cool irrigation lines.
- Calibrate injection pump to evenly distribute Retaliate BN.
- Irrigate for 15-30 minutes prior to starting treatment to cool lines and establish a wetting pattern under plants.
- Add Retaliate BN to container containing no less than 5 gallons of water (temp should be 40-85F), stir contents, let sit for 5 minutes and mix again.
- Empty bucket into injection tank and add sufficient water.
- Irrigate after application with 0.5-1 inch of water.

BEST PRACTICES

- Store nematodes cool: 38°F/3°C until ready for use.
- Do not Subdivide. Once opened, use entire tray immediately.
- Store unopened. Follow product specific shelf life. Do not freeze.
- Allow packages to come to room temperature before mixing.
- Do not apply nematicide within 2 weeks before or after application.
- Avoid direct sunlight. Apply early morning or evening.
- Apply mixed solution as soon as possible. Do not leave nematode/water mixture sitting for more than 2 hours. Not aquatic nematodes.
- Remove all fine filters (<50 mesh).
- Ideal injection pressure is ~200 psi or below.
- Do not exceed 300 psi
- Maintain proper soil moisture for 2 weeks.
- Nematodes have limited horizontal movement, moving vertically in soil until they find a pest. Thorough coverage of the area is important for achieving control.



DISTRIBUTED BY:
SOUTHEAST AG SOLUTIONS, LLC

Water & Soil Biologicals

P.O. Box 7964, Sebring, FL 33872

863.273.2768

www.southeastagsolutions.com



MANUFACTURED BY:

Asymmetry

104 Three West Parkway, Bldg 300
Villa Rica, GA 30180